

Computing eccentric connectivity index of nanostar dendrimers

Sara Mehdipour, Mehdi Alaeiyan, Ali Nejati

Department of Mathematics, Karaj Branch, Islamic Azad University, Karaj, Iran
a.nejati56@yahoo.com

Abstract: Let G be a molecular graph, the *eccentric connectivity index* of G is defined as $\xi^c(G) = \sum_{u \in V(G)} \deg(u) \cdot ecc(u)$, where $\deg(u)$ denotes the degree of vertex u and $ecc(u)$ is the largest distance between u and any other vertex v of G , namely, eccentricity of u . In this study, we present exact expressions for the eccentric connectivity index of two infinite classes of nanostar dendrimers.

Keywords: eccentric connectivity index, nanostar dendrimers, topological index, eccentricity

Introduction

Molecular descriptors are playing significant role in chemistry, pharmacology, etc. Among them, topological indices have a prominent place (Todeschini et al., 2000). There are numerous of topological descriptors that have found some applications in theoretical chemistry, especially in QSPR/QSAR research.

More recently, a new topological index, *eccentric connectivity index*, has been investigated. This topological model has been shown to give a high degree of predictability of pharmaceutical properties, and may provide leads for the development of safe and potent anti-HIV compounds. We encourage readers to consult papers (Dureja et al., 2005, 2006, 2009; Kumar et al., 2006, 2007; Lather et al., 2005; Sardana et al., 2001, 2002; Sharma et al., 1997) for some applications and papers (Morgan et al., 2010; Ilic et al., 2001; Xu X; Zhou et al., 2010; Ashrafi et al., 2011) for the mathematical properties of this topological index.

Dendrimers are highly branched macromolecules. They are being investigated for possible uses in nanotechnology, gene therapy, and other fields. Each dendrimer consists of a multifunctional core molecule with a dendritic wedge attached to each functional site. The core molecule without surrounding dendrons is usually referred to as zeros generation. Each successive repeat unit along all branches forms the next generation, 1st generation and 2nd generation and so on until the terminating generation. The topological study of these macromolecules is the aim of following articles, see (Khoramdel et al., 2008; Ashrafi et al., 2008; Karbasioun et al., 2009; Yousefi-Azari et al., 2008) for details.

Now, we introduce some notation and terminology. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. Let $\deg(u)$ denote the degree of the vertex u in G . If $\deg(u) = 1$, then u is said to be a *pendent*

vertex. An edge incident to a pendent vertex is said to be a *pendent edge*. For two vertices u and v in $V(G)$, we denote by $d(u, v)$ the distance between u and v , i.e., the length of the shortest path connecting u and v . The *eccentricity* of a vertex u in $V(G)$, denoted by $ecc(u)$, is defined to be

$$ecc(u) = \max\{d(u, v) \mid v \in V(G)\}$$

The *diameter* of a graph G is defined to be $\max\{ecc(u) \mid u \in V(G)\}$. The *eccentric connectivity index*, $\xi^c(G)$, of a graph G is defined as

$$\xi^c(G) = \sum_{u \in V(G)} \deg(u) \cdot ecc(u)$$

where $\deg(u)$ is the the degree of a vertex u and $ecc(u)$ is it's eccentricity.

The second author of this paper in some joint works computed some topological indices of some molecular graphs related to eccentricity (Alaeiyan et al., 2013, 2015; Nejati et al., 2014, 2015). Yarahmadi investigated some topological indices of nanostar dendrimers in (Yarahmadi et al., 2010, 2011) and Ashrafi. A. R. in (Ashrafi et al., 2012) calculated eccentric connectivity index of a class of nanostar dendrimers. Also in (Nilanjan et al., 2016) Nilanjan presented exact expressions for the F-index and F-polynomial of six infinite classes of nanostar dendrimers. In this paper we computed the eccentric connectivity index of two infinite classes of nanostar dendrimers. Their structures are given in Figs. 1, 3.

Results

Suppose D_n denotes the molecular graph of a dendrimer with exactly generations depicted in Fig. 1 where $n \geq 1$. The total number of vertices and edges of D_n are calculated as $57 \times 2^{n-1} - 38$ and $33 \times 2^n - 45$ respectively. In the following theorem we calculate the eccentric connectivity index of D_n .

Theorem 1. The eccentric connectivity index of D_n is computed as

$$\xi^c(D_n) = (1188 \times 2^n - 810)n - 1485 \times 2^n + 1665.$$




Fig. 1. The nanostar dendrimer $D_n(n=2)$.




Fig. 2. The eccentricities in a third of D_2 .

Proof . Considering Fig. 2 and Table. 1, it can be seen that, we have $9n-4$ types of vertices in D_n , based on their eccentricities. We have $3 \times 2^{n-1}$ numbers of vertices of type 1 with maximum eccentric connectivity equals to $18n-10$ (red vertices). The number of vertices of type 2, 3 is 3×2^n and their eccentricities are $18n-11$ and $18n-12$ respectively. Also we have $3 \times 2^{n-1}$ numbers of vertices of type 4 with eccentric connectivity equals to $18n-13$ and so it continues until we have three vertices of type $9n-5$ with eccentric connectivity equals to $9n-4$ and finally there is a vertex of type $9n-4$ with minimum eccentric connectivity equals to $9n-5$ (blue vertex). It is clear that for any vertex u in D_n , $\deg(u) = 2$ or $\deg(u) = 3$ (see Table. 1). Therefore we have

$$\begin{aligned} \xi^c(D_n) &= \sum_{u \in V(G)} \deg(u) \cdot ecc(u) = \\ &= 2 \times 3 \times 2^{n-1} \times (18n-10) \\ &+ 3 \times 2^n \times 2 \times (18n-11) + 3 \times 2^n \times 2 \times (18n-12) \\ &+ 3 \times 2^{n-1} \times 3 \times (18n-13) + 3 \times 1 \times (9n-5) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-2-k} \times 3 \times (18n-14-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-2-k} \times 3 \times (18n-15-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-1-k} \times 2 \times (18n-16-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-1-k} \times 2 \times (18n-17-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-2-k} \times 3 \times (18n-18-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-2-k} \times 3 \times (18n-19-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-1-k} \times 2 \times (18n-20-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-1-k} \times 2 \times (18n-21-9k)) \\ &+ \sum_{k=0}^{n-2} (3 \times 2^{n-2-k} \times 3 \times (18n-22-9k)) \\ &= 6 \times 2^{n-1} \times (18n-10) + 6 \times 2^n \times (18n-11) \\ &+ 6 \times 2^n \times (18n-12) + 9 \times 2^{n-1} \times (18n-13) \\ &+ 3 \times (9n-5) + \sum_{k=0}^{n-2} [9 \times 2^{n-2-k} \times (90n-45k-80) \\ &+ 6 \times 2^{n-1-k} \times (72n-36k-74)]. \end{aligned}$$

Now with simplification in MATLAB software (Matlab et al., 2012) we have

$$\xi^c(D_n) = (1188 \times 2^n - 810)n - 1485 \times 2^n + 1665.$$

Then this proof is completed.

Now we consider another class of dendrimer, denoted as $NS_5(n)$ denotes the molecular graph of a dendrimer with exactly n generations depicted in Fig. 3 where $n \geq 1$.

In the following theorem we calculate the eccentric connectivity index of $NS_5(n)$.

Theorem 2. The eccentric connectivity index of $NS_5(n)$ is computed as

$$\xi^c(NS_5(n)) = (1200 \times 2^n + 270)n + 2544 \times 2^n + 1275.$$

Proof . Considering Fig. 4 and Table. 2, it can be seen that, we have $5n+14$ types of vertices in

Table 1. Types of vertices in D_n .

Types of Vertices	Num	$ecc(u)$	$deg(u)$
1	$2^{n-1} \times 3$	$18n - 10$	2
2	$2^n \times 3$	$18n - 11$	2
3	$2^n \times 3$	$18n - 12$	2
4	$2^{n-1} \times 3$	$18n - 13$	3
5	$2^{n-2} \times 3$	$18n - 14$	3
6	$2^{n-2} \times 3$	$18n - 15$	3
...
$9n - 6$	6	$9n - 3$	2
$9n - 5$	3	$9n - 4$	3
$9n - 4$	1	$9n - 5$	3

$NS_5(n)$, based on their eccentricities. We have 9×2^n numbers of vertices of type 1 with maximum eccentric connectivity equals to $10n+28$ (red vertices). Also we have 3×2^n numbers of vertices of types 2, 3 with eccentric connectivity equals to $10n+27$ and $10n+26$ respectively. The number of vertices of type 4 is 3×2^n and their eccentricity is $10n+25$ and so it continues until we have six vertices of type $5n+13$ with eccentric connectivity equals to $5n+16$ and finally there are three vertices of type $5n+14$ with minimum eccentric connectivity equals to $5n+15$ (blue vertices). It is clear that for any vertex u in $NS_5(n)$, $deg(u) = 1$ or $deg(u) = 2$ or $deg(u) = 3$. (See Table. 2). Therefore we have




Fig. 3 Polymer dendrimer $NS_5(n)(n = 2)$.




Fig. 4 The eccentricities in a third of $NS_5(2)$.

$$\begin{aligned}
 \xi^c(NS_5(n)) &= \\
 &= \sum_{u \in V(G)} \deg(u) \cdot ecc(u) = 3 \times 2^{n+1} \times (10n + 28) \\
 &+ 3 \times 2^n \times (2) \times (10n + 28) + \sum_{k=1}^{n-1} \frac{3 \times 2^{n+1}}{2^{k-1}} (3)(10n - 5k + 32) \\
 &+ \sum_{k=1}^{n-1} \frac{3 \times 2^{n+1}}{2^{k-1}} (2)(10n - 5k + 31) + \\
 &+ \sum_{k=1}^{n-1} \frac{3 \times 2^n}{2^{k-1}} (3)(10n - 5k + 30) \\
 &+ \sum_{k=1}^{n-1} \frac{3 \times 2^n}{2^{k-1}} (2)(10n - 5k + 29) + \\
 &+ \sum_{k=1}^{n-1} \frac{9 \times 2^{n-1}}{2^{k-1}} (2)(10n - 5k + 28) \\
 &+ \sum_{k=1}^2 (4-k)(12)(5n-k+33) + \sum_{k=1}^3 (6)(2)(5n-k+30) \\
 &+ \sum_{k=1}^2 (6)(3)(5n-4k+34) + \sum_{k=1}^2 (2)(3k)(5n-3k+29) \\
 &+ \sum_{k=1}^2 (6)(3)(5n-k+26) + \sum_{k=1}^2 (3)(3)(5n-k+23) \\
 &+ \sum_{k=1}^2 (2)(6)(5n-k+21) + \sum_{k=1}^2 (3)(3)(5n-2k+20) \\
 &+ \sum_{k=1}^2 (2)(3)(5n-k+18) + \\
 &+ (3)(1)(5n+17) + (3)(3)(5n+15)
 \end{aligned}$$

$$\begin{aligned}
 &= 12 \times 2^n \times (10n + 28) \\
 &+ \sum_{k=1}^{n-1} \left[\frac{2^{n+1}}{2^{k-1}} (150n - 75k + 476) + \right. \\
 &+ \frac{2^n}{2^{k-1}} (150n - 75k + 444) \\
 &+ \left. \frac{2^{n-1}}{2^{k-1}} (180n - 90k + 504) \right] + \sum_{k=1}^3 12(5n - k + 30) \\
 &+ \sum_{k=1}^2 (600n - 405k - 30kn - 6k^2 + 3411) + 3 \times (5n + 17) \\
 &+ 9 \times (5n + 15).
 \end{aligned}$$

Now with simplification in MATLAB software (Matlab et al., 2012) we have

$$\xi^c(NS_5(n)) = (1200 \times 2^n + 270)n + 2544 \times 2^n + 1275.$$

Then this proof is completed.

Table 2. Types of vertices in $NS_5(n)$.

Types of Vertices	Num	$ecc(u)$	$deg(u)$
1	$3 \times 2^{n+1}$	$10n + 28$	1
1	3×2^n	$10n + 28$	2
2	$3 \times 2^{n+1}$	$10n + 27$	3
3	$3 \times 2^{n+1}$	$10n + 26$	2
4	3×2^n	$10n + 25$	3
5	3×2^n	$10n + 24$	2
...
$5n + 11$	3	$5n + 18$	3
$5n + 12$	3	$5n + 17$	2
$5n + 12$	3	$5n + 17$	1
$5n + 13$	3	$5n + 16$	3
$5n + 13$	3	$5n + 16$	2
$5n + 14$	3	$5n + 15$	3

Conclusions

The eccentric connectivity index has been employed successfully for the development of numerous mathematical models for the prediction of biological activities of diverse nature. Dendrimers are large and complex molecules with well tailored chemical structures. In this paper, exact formulas for the eccentric connectivity index of two infinite classes of nanostar dendrimers are given. It would be interesting for future study to investigate other topological indices such as edge version of eccentric connectivity index of these nanostar dendrimers.

References

- Alaeiyan M, Asadpour J and Mojarad R (2013) A Numerical Method for MEC Polynomial and MEC Index of One-Pentagonal Carbon Nanocones, Fullerenes, Nanotubes and Carbon Nanostructures. 21: 825–835.
- Alaeiyan M, Asadpour J and Mojarad R (2013) A Study of an Infinite Family of Carbon Nanocones by the Eccentricity Connectivity polynomial, Fullerenes, Nanotubes and Carbon Nanostructures. 21: 849–860.
- Alaeiyan M, Asadpour J and Mojarad R (2015) A new method for computing eccentric connectivity polynomial of an infinite family of linear polycene parallelogram benzenoid, *optoelectronics and advanced materials – rapid communications*. 9: 813–815.
- Ashrafi AR, Doslic T and Saheli M (2011) The eccentric connectivity index of $TUC4C8(R)$ nanotubes, *MATCH Commun. Math. Comput. Chem.* 65: 221–230.
- Ashrafi AR, Mirzargar M (2008) Indian J. Chem. 47A: 538.
- Ashrafi AR, Saheli M (2012) COMPUTING eccentric connectivity index of a class of nanostar dendrimers, Kragujevac J. Sci. 34: 65–70.
- Dureja H, Madan AK (2005) Topochemical models for prediction of cyclindependent kinase 2 inhibitory activity of indole-2-ones, *J. Mol. Model.* 11: 525–531.
- Dureja H, Madan AK (2006) Topochemical models for the prediction of permeability through blood-brain barrier, *Int. J. Pharm.* 323: 27–33.
- Dureja H, Madan AK (2009) Predicting anti-HIV activity of dimethylaminopyridin-2-ones: Computational approach using topochemical descriptors, *Chem. Biol. Drug Des.* 73: 258–270.
- Ilic A, Gutman I (2011) Eccentric connectivity index of chemical trees, *MATCH Commun. Math. Comput. Chem.* 65: 731–744.
- Karbasioun A, Ashrafi AR (2009) Maced. J. Chem. Eng. 28: 49.
- Khoramdel MH, Yousefi-Azari H, Ashrafi AR (2008) Indian J. Chem. 47A: 1503.
- Kumar V, Madan AK (2006) Application of graph theory: Prediction of cytosolic phospholipase A(2) inhibitory activity of propan-2-ones, *J. Math. Chem.* 39: 511–521.
- Kumar V, Madan AK, Application of graph theory: Models for prediction of carbonic anhydrase inhibitory activity of sulfonamides, *J. Math. Chem.*, (2007), 42: 925–940.
- Lather V, Madan AK (2005) Application of graph theory: Topological models for prediction of CDK-inhibitory activity of aloisines, *Croat. Chem. Acta.* 78: 55–61.
- MATLAB and Statistics Toolbox Release 2012b, The MathWorks, Inc., Natick, Massachusetts, United States.
- Morgan MJ, Mukwembi S, Swart HC (2010) On the eccentric connectivity index of a graph, *Disc. Math.* doi:10.1016/j.disc.12.013.
- Nejati A, Alaeiyan M (2014) The edge version of MEC index of one-pentagonal carbon nanocones, *Bulgarian Chemical Communications.* 46: 462–464.
- Nejati A, Alaeiyan M (2015) The edge version of MEC index of linear polycene parallelogram benzenoid, *optoelectronics and advanced materials – rapid communications*. 9: 813–815.
- Nilanjan De, Abu Nayeem SkMd (2016) Computing the F-index of nanostar dendrimers. Pacific Science Review A: Natural Science and Engineering. 18: 14–21.
- Sardana S, Madan AK (2001) Application of graph theory: Relationship of molecular connectivity index, Wiener index and eccentric connectivity index with diuretic activity, *MATCH Commun. Math. Comput. Chem.* 43: 85–98.
- Sardana S, Madan AK (2002) Application of graph theory: Relationship of antimycobacterial activity of quinolone derivatives with eccentric connectivity index and Zagreb group parameters, *MATCH Commun. Math. Comput. Chem.* 45: 35–53.
- Sharma V, Goswami R, Madan AK (1997) Eccentric connectivity index: A novel highly discriminating topological descriptor for structure-property and structure-activity studies, *J. Chem. Inf. Model.* 37: 273–282.
- Todeschini R, Consonni V (2000) Hand book of Molecular Descriptors (Wiley-VCH), Weinheim.
- Xu X, The eccentric connectivity index of trees of given order and matching number, submitted for publication.
- Yarahmadi Z, Fathtabar GH (2011) The Wiener, Szeged, PI, vertex PI, the first and second Zagreb indices of N-branched phenylacetylenes dendrimer, *MATCH Commun. Math. Comput. Chem.* 65(1): 201–208.
- Yarahmadi Z (2010) Eccentric Connectivity and Augmented Eccentric Connectivity Indices of N-Branched Phenylacetylenes Nanostar Dendrimers, *Iranian Journal of Mathematical Chemistry.* 1: 105–110.
- Yousefi-Azari H, Ashrafi AR, Bahrami A, Yazdani J (2008) Asian. J. Chem. 20: 15.
- Zhou B (2010) On eccentric connectivity index, *MATCH Commun. Math. Comput. Chem.* 63: 181–198.